

LECTURE 9: DEGREE LOWER BOUNDS FOR POLYNOMIAL CALCULUS

Fix field $\mathbb{F}$ variables $\vec{x} = \{x_1, \dots, x_n\}$

Work in $\mathbb{F}[\vec{x}] / \{x_j^2 - x_j \mid j \in [n]\}$

so all polynomials multilinear

Also throw in variables $\bar{x}_j$ when convenient — won't need them today

POLYNOMIAL CALCULUS DERIVATION (sequence of polynomials $\pi = (p_1, p_2, \dots, p_k)$ such that

p_i is

INPUT POLYNOMIAL

$p_i \in \mathcal{P}$

BOOLEAN AXIOM

$$x_j^2 - x_j$$

except we don't need these in multilinearity

COMPLEMENTARITY AXIOM

$$x_j + \bar{x}_j - 1$$

or derived from $p_j, p_k \in \pi$, $j, k < i$ by

LINEAR COMBINATION

$$\frac{\alpha q + \beta r}{\alpha q + \beta r}$$

$\alpha, \beta \in \mathbb{F}$

MULTIPLICATION

$$\frac{q}{mq}$$

m monomial

A POLYNOMIAL CALCULUS REFUTATION of set of polynomials $\mathcal{P}$ is derivation from $\mathcal{P}$ ending in $p_k = \underline{1}$

Size of p $S(p) = \#$ monomials in representation of p as linear combination of monomials

$$S(\pi) = \sum_i S(p_i)$$

$$\text{Deg}(\pi) = \max \text{ total degree of any polynomial } p_i$$

$$\text{Spec}(P \perp) = \exp\left(-\Omega\left(\frac{(\text{Deg}(P \perp) - \text{Deg}(P))^2}{|\text{Vars}(P)|}\right)\right)$$

Focus on proving degree lower bounds

Build generalized constraint-variable incidence graph (CVIG)

sets of polynomials P over variables V

Partition $P = Q \cup \bigcup_{i=1}^m P_i$

Divide $V = \bigcup_{i=1}^n V_i$

OVERLAP $ol(V) = \max_x |\{V_i \mid x \in V_i\}|$

$(U, V)_Q$ -graph

Bipartite graph with

- left vertex set $U = \{P_1, P_2, \dots, P_m\}$
- right vertex set $V = \{V_1, \dots, V_n\}$
- Edge (P, V) if $\text{Vars}(P) \cap V \neq \emptyset$

The edge (P, V) is RESPECTFUL or PC if satisfies wins following game on (P, V)

- (1) Satisfies commits to $\rho: V \rightarrow \{T, \perp\}$ such that if for $q \in Q$ it holds that $\text{Vars}(q) \cap V \neq \emptyset$, then ρ satisfies q
- (2) Adversary plays α s.t. $\alpha(Q) = T$
- (3) Define $\alpha'(x)$ by

$$\alpha'(x) = \begin{cases} \rho(x) & \text{if } x \in V \\ \alpha(x) & \text{otherwise} \end{cases}$$

Satisfies wins if $\alpha'(P \wedge Q) = T$

Recall for $G = (U \cup V, E)$, $U' \subseteq U$

$$\partial(U') = \{v \in V : |N(v) \cap U'| = 1\}$$

BOUNDARY or UNIQUE NEIGHBOURS of U'

PC BOUNDARY of U' in $(U, V)_Q$ -graph

$$\partial_Q^{PC}(U') = \{v \in V \mid v \in \partial(U') \text{ and for } P_v \in N(v) \cap U' \text{ } (P_v, v) \text{ is a PC edge}\}$$

$(U, V)_Q$ -graph is (s, δ, ξ, Q) -PC EXPANDER

if $\forall U' \subseteq U, |U'| \leq s$ it holds that
with overlap ℓ if $ol(V') \leq \ell$

$$|\partial_Q^{PC}(U')| = \delta |U'| - \xi$$

THEOREM 2 [MN '24]

Suppose $\forall p \in \mathcal{P} \quad |Vars(p)| \leq \frac{\delta s - 2\xi}{2\ell}$

$(U, V)_Q$ (s, δ, ξ, Q) -expander with overlap ℓ

$\forall U', |U'| \leq s \quad U' \cup Q$ satisfiable.

Then

$$Deg(\mathcal{P} + \perp) > \frac{\delta s - 2\xi}{2\ell}$$

COROLLARY 3 [MN '24]

Suppose that $\forall p \in \mathcal{P} \quad |Vars(p)| \leq \frac{\delta s}{2\ell}$

and $(U, V)_Q$ -graph is $(s, \delta, 0, Q)$ -PC expander (i.e., $\xi = 0$)

Then

$$Deg(\mathcal{P} + \perp) > \frac{\delta s}{2\ell}$$

How can we use this to prove lower bounds?

Build $(\mathcal{U}, \mathcal{V})_Q$ -graphs and show that they are PC expanders.

THEOREM 4 [Alkhnovich & Razborov '03]

Let F be a CNF formula such that the clause-variable incidence graph $G(F)$ is an (s, δ) -boundary expander for some $\delta > 0$. Then $\text{Deg}(F \vdash 1) > \delta s/2$.

COROLLARY 5

for $k \geq 3$
(with $\Theta(n)$ clauses over n variables)

Random k -CNF formulas require polynomial calculus proofs of exponential size $\exp(\Omega(n))$ asymptotically almost surely (and lineardegree $\Omega(n)$)

Proof of Cor 5 The CVIG of such a random formula is a random k -left-regular bipartite graph, which is an excellent expander a.a.s.

Proof of Thm 4 Axiom clauses/polyomials have $k = O(1)$ variables. Set $Q = \emptyset$, $\mathcal{U} = \text{singleton set of clauses}$, $\mathcal{V} = \text{singleton sets of variables}$. All edges $\stackrel{(C, x)}{\text{are}}$ PC good - can assign x to fix $C = T$. $\square$

GRAPH ORDERING PRINCIPLE $\text{GOP}(G)$

"There exists a totally ordered set on n elements without a minimal element."

Variables $x_{u,v} = \text{"element } u < \text{element } v \text{"}$
 $u, v \in [n]$

Bounded-degree graph $G = (V, E)$ $V = [n]$

DEF Non-bipartite boundary expander

Graph $G = (V, E)$ is an (s, δ) -BOUNDARY EXPANDER if $\forall V' \subseteq V, |V'| \leq s$, it holds that $|\partial(V')| \geq \delta |V'|$, where

$$\partial(V') = \{v \in V \setminus V' : |N(v) \cap V'| = 1\}$$

$$\bar{x}_{u,v} \vee \bar{x}_{v,w} \vee x_{u,w}$$

$$u, v, w \in V \\ u \neq v \neq w \neq u$$

TRANSITIVITY

$$\bar{x}_{u,v} \vee \bar{x}_{v,u}$$

$$u, v \in V \\ u \neq v$$

ASYMMETRY

$$x_{u,v} \vee x_{v,u}$$

$$u, v \in V \\ u \neq v$$

TOTALITY

$$\bigvee_{u \in N(v)} x_{u,v}$$

$$v \in V$$

NON-MINIMALITY

"v is not minimal as witnessed by some neighbour u of v w.r.t. G"

For $G = K_n$, this formula is known as the LINEAR ORDERING PRINCIPLE FORMULA (or LEAST NUMBER PRINCIPLE FORMULA or GRAPH TAUTOLOGY)

THEOREM 6 [Galexi & Lauria '10]

For any non-bipartite (s, δ) -boundary expander G it holds that

$$\text{Deg}(GOP(G) \vdash \perp) > \frac{\delta s}{4}$$

Remark 1 Result in Galexi-Lauria has slightly better constants — we optimize for simplicity of exposition (GL'10 gets lower bounds for 3-col here we need 5-col)

Remark 2 Random δ -regular graphs are

$(\Omega(n), \delta > 0)$ -boundary expanders a.a.s. — calculation

Remark 3 Thm 6 yields 5-CNF formulas over

$\Theta(n^2)$ variables with repetition degree $\Omega(n)$
 [Spilmarck '96] showed that there are short resolution proofs. ^{$O(n^3)$} So these formulas demonstrate that size-width/degree claims in [IPS '99] and [BW '01] are essentially tight. Might look at Spilmarck's resolution proof later in the course

Proof of Thm 6 Let $\mathcal{Q}$ consist of all transitivity, asymmetry, and totality axioms (enforcing that assignments are total orders)

Left vertex sets $\mathcal{U}$ = singleton sets with non-minimality axioms

Right vertex sets $\mathcal{V} = \{V_v \mid v \in V\}$ where

$$V_v = \{x_{u,w} \mid u=v \text{ or } w=v\}$$

set of all variables that mention v .

Every variable $x_{u,w}$ occurs in V_u and V_w , so overlap = 2.

We claim that all edges (P_v, V_w) are PC-good.
 $w \in \{v\} \cup N(v)$

If $w \in N(v)$, then let

$$\begin{aligned} f_w(x_{w,u}) &= \top && \text{for all } u \text{ such that } x_{u,w} \text{ or } x_{w,u} \text{ are in } V_w \\ f_w(x_{u,w}) &= \perp \end{aligned}$$

i.e., f_w sets w to be locally smallest element.

$P_v = \{ \bigvee_{u \in N(v)} x_{u,v} \}$ is satisfied.

For Q , asymmetry and totality axioms are either untouched or satisfied, since if g assigns $x_{u,w}$ it also assigns $x_{w,u}$ and exactly one of the variables is true and the other false.

$$\overline{x_{a,b}} \vee \overline{x_{b,c}} \vee x_{a,c}$$

For transitivity if $g(x_{a,c}) = T$, then we're OK and if $g(x_{a,c}) = \perp$ then $c = w$ and we also have $g(x_{b,c}) = \perp$

$a = w$ and

Similarly, if $g(x_{ab}) = T$ then $g(x_{a,c}) = T$
if $g(x_{b,c}) = T$ then $b = w$ and $g(x_{ab}) = \perp$

Consider vertex set $V' \subseteq V(G)$. By discussion above, we have

$$\partial_Q(\cup_{v' \in V'} \{P_{v'}\}) \supseteq \{V_w \mid w \in \partial(V')\}$$

in $(\mathcal{U}, V)_Q$ -graphs, so the $(\mathcal{U}, V)_Q$ -graph has at least as good expansion parameters.

We get

$$\text{Deg}(G \cup \{G \cup \{G\} + 1) > \frac{\delta S}{2L} = \frac{\delta S}{4} \quad \square$$

If $w = v$ then let

$$g_v(x_{v,u}) = \perp$$

$$g_v(x_{u,v}) = T$$

i.e., g_v sets v to locally largest element

Exactly same argument shows that Q is OK

$$G = (U \cup V, E)$$

Pigeon u can fly to holes $N(u)$

$$\forall v \in N(u) \quad x_{u,v}$$

$$u \in U$$

PIGEON AXIOMS

$$\bar{x}_{u,v} \vee \bar{x}_{u',v}$$

$$\begin{aligned} &v \in V \\ &u, u' \in N(v) \\ &u \neq u' \end{aligned}$$

HOLE AXIOMS

$$\bar{x}_{u,v} \vee \bar{x}_{u,v'}$$

$$\begin{aligned} &u \in U \\ &v \neq v' \in N(u) \end{aligned}$$

FUNCTIONALITY AXIOMS

$$\forall u \in N(v) \quad x_{u,v}$$

$$v \in V$$

ONTO AXIOMS

Standard pigeonhole principle formulas

for graph $K_{n+1, n}$ by restriction

Get formula over G

$$f_G(x_{u,v}) = \begin{cases} 0 & \text{if } (u,v) \notin E \\ * & \text{otherwise} \end{cases}$$

So lower bounds for graph PHP formulas imply lower bounds for standard PHP formulas

Let $\text{Onto-PHP}(G)$ contain pigeon, hole, and onto axioms

$\vdash \text{FPHP}(G) \vdash \text{Onto-PHP}(G)$ pigeon, hole, and functionality axioms

THEOREM 7 (Essentially [AR'03])

Let $G = (U \cup V, E)$ be an (s, δ) -boundary expander without isolated vertices in V . Then

$$\text{Deg}(\text{Onto-PHP}(G) \vdash \perp) \geq \frac{s\delta}{2}.$$

Proof sketch Let $Q =$ onto & hole axioms

Let $U =$ singleton sets with pigeon axioms.

Let $V_v = \{x_{u,v} \mid u \in N(v)\}$, i.e., variables are partitioned w.r.t. holes. All edges are PC-good, and the $(U, V)_Q$ -graph is isomorphic to G . (Note, though, that pigeons might be sent to several holes to satisfy onto axioms in Q .) $\square$

This does not work for $FPHP(G)$, if we try to substitute functionality axioms for onto axioms in Q .

For edge (P_u, V_v) , need to set $x_{u,v} = T$

But $(\bar{x}_{u,v} \vee \bar{x}_{u,v'}) \upharpoonright_{x_{u,v}} = \bar{x}_{u,v'}$ — clause touched but not satisfied

THEOREM 8 [MN'24]

Let $G = (U \cup V, E)$ be a bipartite (s, δ) -^{boundary}expander with left degree $\leq d$. Then

$$\text{Deg}(FPHP(G) \vdash \perp) > \frac{\delta s}{2cd}$$

Proof Construct $(U, V)_Q$ -graph as follows.

$Q =$ hole & functionality axioms

$U =$ singleton sets with pigeon axioms

$V = \{V_v \mid v \in V\}$ where

$$V_v = \{x_{u',v'} \mid u' \in N(v) \text{ and } v' \in N(u')\}$$

- i.e.
- start with hole v
 - go to all pigeons u' that can fly to v
 - consider all holes v' such pigeons u' can fly to

Overlap is $\leq d$, since $x_{u,v}$ can only appear PCD in sets $V_{v'}$ such that $v' \in N(u)$, and $|N(u)| \leq \text{left degree } d$.

G and $(\mathcal{U}, \mathcal{V})_Q$ is same graph

If $(u, v) \in E(G)$, then clearly $\exists$ edge $P_u \leftrightarrow V_v$

Suppose (P_u, V_v) is edge in $(\mathcal{U}, \mathcal{V})_Q$

Then by construction $u \in N(v)$, so $(u, v) \in E(G)$.

Finally, we claim that all edges are PC good.

Consider $P_u = \{ V_{w \in N(u)} x_{u,w} \}$ and V_v s.t. $v \in N(u)$, i.e., (P_u, V_v) is edge. Choose

$$g_{u,v}(x_{u,v'}) = \begin{cases} \top & \text{if } u'=u \text{ \& } v'=v \\ \perp & \text{otherwise} \end{cases}$$

$V_{w \in N(u)} x_{u,w}$ is satisfied

Hole axioms for v satisfied, since $g(x_{u,v}) = \top$

All other hole axioms untouched or satisfied, since all other variables in V_v set to $\perp$.

Functionality axioms for u satisfied,

since $g(x_{u,v'}) = \perp$

All other functionality axioms either untouched or satisfied, since all other variables in V_v set to $\perp$.

Same graph and all edges good $\rightarrow$ same expansion

Overlap $\ell \leq d$ yields degree lower bound $\frac{5\epsilon}{(2d)}$



$$S_{PCR}(\text{FPHP}_n^{n+1} \vdash \perp) = \exp(-\Omega(n))$$

Proof Apply restriction to $K_{n+1, n}$ to get, say, δ -left-regular $(\Omega(n), \delta > 0)$ -boundary expander G .

Theorem 8 $\Rightarrow$ degree lower bound $\Omega(n)$ for $\text{FPHP}(G)$

[IPS'99] $\Rightarrow$ exponential size lower bound for $\text{FPHP}(G)$

Since $\text{FPHP}_n^{n+1} \upharpoonright_G = \text{FPHP}(G)$ and restrictions preserve refutations, means that same lower bound applies for FPHP_n^{n+1} $\square$

Some questions

Results for resolution
in [BIS'07]

- ① Can we get PC lower bounds for graph coloring?
Yes, future lecture
- ② What about clique or vertex cover?
- ③ PC lower bounds for Ramsey theory problems?
- ④ Degree lower bounds for dense linear order formulas
Resolution
[Atserias, Dalmau '08]
- Resolution results
[Carlucci, Guler, Lauria '16]
[Lauria, Pudlák, Rödl, Thapen '17]
[Pudlák '12]
- ⑤ PC size lower bounds for weak pigeonhole principle (WPHP) formulas with $\geq n^2$ pigeons
Understood for resolution Raz & Razborov
Also for graph-WPHP [de Rezende, Nordström, Risse, Solovov '25]

The lower bound method we have presented works regardless of the field. What about lower bounds that depend on the field?

(6) [Alekhnovich & Razborov '03] have a version of their method that depends on the field. Can it be combined with generalized CVIGs and filtering sets $\mathcal{A}$?

(7) What about Ono-FPHP formulas in fields $\mathbb{F}$ of characteristic p with $n + p^d$ pigeons and n holes? Upper & lower bounds for Nullstellensatz in [Beame & Rieš '98]
Nothing known for PC

(8) What about perfect matching on sparse non-bipartite graphs G with $|V(G)|$ odd? [Austin & Rieš '22] Degree lower bound $\Omega(\frac{n}{\log n})$ for random d -regular graphs.


Issues: (a) d has to be very large
(b) log factor loss

Proof by reduction to Tseitin formulas.
Direct argument would be desirable

PROOF STRATEGY FOR DEGREE LOWER BOUNDS

PCD XIII

Give our approximation of what
can be derived in degree D by
defining operator R
such that if p derivable
from $\mathcal{P}$ in degree $\leq D$,
then $R(p) = 0$



Then show $R(1) \neq 0$

/// derivable in degree D

/// $\{p \mid R(p) = 0\}$

LEMMA 10 [Razborov '98]

Let $\mathcal{P}$ set of multilinear polynomials, $D \in \mathbb{N}^+$

Suppose $\exists$ linear operator R on multilinear
polynomials over $\text{Vars}(\mathcal{P})$ of degree $\leq D$ such that

(1) $R(f) = 0$ for all axioms $f \in \mathcal{P}$

(2) For every term t with $\text{Deg}(t) < D$ and
every variable x it holds that $R(xt) = R(x)R(t)$

(3) $R(1) \neq 0$

Then $\text{Deg}_{\text{PC}}(\mathcal{P} \vdash 1) > D$

(or $> D/2$ if
we want to
play it safe)

Proof By forward induction over derivation Π .
Change multiplication rule to only multiply
one variable at a time.

VARIABLE
MULTIPLICATION $\frac{p}{xp}$

Modified multiplication
rule

Will prove degree lower bound in this setting
Strictly speaking, maybe lose factor 2 for
monomial multiplication at get degree LB $> D/2$

Or work slightly harder on property (2) PCD XIV
to get it for monomial multiplication. Might be possible.
Will not worry about this

~~~~~ DETOUR ~~~~~

What is the issue?

Could have  $p = \sum_i \alpha_i m_i$

Multiply by  $t$  to get  $tp = \sum_i \alpha_i t m_i$

Could have high-degree terms  $t m_1$  &  $t m_2$  that cancel

But maybe we don't have cancellation if  
have to multiply by  $x \in \text{Vars}(t)$  one at a time  
Variable multiplication can only double  
the degree

$p$  has degree  $D_1 \geq \max \{ \text{Deg}(m_1), \text{Deg}(m_2) \}$   
 $tp$  has degree  $D_2 \geq \text{Deg}(t)$

$\forall i \text{ Deg}(t m_i) \leq D_1 + D_2 \leq 2 \max \{ D_1, D_2 \}$

By induction over  $\pi = (p_1, p_2, \dots, p_k)$

$R(p_i) = 0$

Base case:  $p_i \in P$   $R(p_i) = 0$  by property (1)

Inductive step  $p_i$  derived by  $\frac{q}{\alpha q + \beta r}$

$$\begin{aligned} R(\alpha q + \beta r) &= \alpha R(q) + \beta R(r) \quad [\text{by linearity}] \\ &= \alpha \cdot 0 + \beta \cdot 0 \quad [\text{by IH}] \\ &= 0 \end{aligned}$$

$p_i$  derived by  $\frac{p}{x^p}$

$p = \sum_i t_i$  for terms  $t_i = x_i \cdot m_i$

$$\begin{aligned}
 R(xp) &= R(x \sum_i t_i) \\
 &= R(\sum_i x t_i) && [\text{linearity}] \\
 &= \sum_i R(x t_i) \\
 &= \sum_i R(x R(t_i)) && [\text{property (2)}] \\
 &= \sum_i R(x R(p)) && [\text{linearity}] \\
 &= \sum_i R(x \cdot 0) && [1\#] \\
 &= 0
 \end{aligned}$$

So if such operator  $R$  exists, all polynomials  $p$  derivable in degree  $\leq D_{\text{axe}} \neq 1$   $\square$

How to construct such an  $R$ ?

$R$  looks a lot like reduction modulo polynomial ideal  $I$

Ideal  $\langle P \rangle$  generated by <sup>input</sup> polynomials  $P$  satisfy (1) and (2)

But not (3), since  $P$  is unsatisfiable iff  $1 \in \langle P \rangle$

Solution Reduce different polynomials modulo different subsets of polynomials of  $P$  that are satisfiable. Piece together so that it looks like real polynomial ideal in degree  $\leq D$

## QUICK REVIEW OF ALGEBRA BASICS

ALG I

Total ordering  $<$  of multivariate monomials over some fixed set of variables is ADMISSIBLE if:

- (a) If  $\text{Deg}(m_1) < \text{Deg}(m_2)$ , then  $m_1 < m_2$
- (b) For monomials  $m_1, m_2, m$  such that  $\text{Vars}(m) \cap (\text{Vars}(m_1) \cup \text{Vars}(m_2)) = \emptyset$ .

and  $m_1 < m_2$  it holds that  $m m_1 < m m_2$

Write  $m_1 \leq m_2$  for  $m_1 < m_2$  or  $m_1 = m_2$

Terms  $t_1 = \alpha_1 m_1$  and  $t_2 = \alpha_2 m_2$  ( $\alpha_i \in \mathbb{F}$ ) are ordered as underlying monomials  $m_1$  &  $m_2$

\*) Tailor-made definition of admissible for our purposes — general definition is, well, more general.

Exact choice of order almost doesn't matter

For concreteness, let us order first w.r.t. degree and then lexicographically  $x_1 \leq x_2 \leq x_3 \leq \dots \leq x_n$

$$x_2 x_3 x_4 < x_2 x_3 x_5 < x_1 x_2 x_3 x_4$$

In what follows write polynomials

$p = \sum_i \epsilon_i$  as sums of terms over distinct monomials.

LEADING TERM  $\text{LT}(p)$  of  $p = \sum_i \epsilon_i$  is largest term  $t_i$  according to  $<$ .

Let  $I$  be ideal in  $\mathbb{F}[\vec{x}] / \{x_j^2 - x_j \mid j \in [n]\}$  ALG II  
ring of multilinear polynomials

Term  $t$  is REDUCIBLE MODULO  $I$  if  $\exists g \in I$   
s.t.  $t = \text{LT}(g)$  and IRREDUCIBLE otherwise.

FACT A Let  $I$  ideal over and  $p$  polynomial  
in  $\mathbb{F}[\vec{x}] / \{x_j^2 - x_j \mid j \in [n]\}$ . Then  $p$   
can be written uniquely as  
$$p = g + r$$
  
for  $g \in I$  and  $r$  sum of irreducible  
terms mod  $I$

Proof  $p$  can be written as  $p = g + r$  for  
 $g \in I$  and  $r$  sum of irreducibles in some  
way by induction over  $\text{LT}(p)$ .

- (i) If  $\text{LT}(p)$  irreducible, then apply IH to  
 $p' = p - \text{LT}(p)$  which has smaller leading term  
(ii) If  $\text{LT}(p)$  reducible, choose  $g \in I$  s.t.  $\text{LT}(g) = \text{LT}(p)$   
and apply IH to  $p' = p - g$ .

In both cases  $p' = g' + r'$  by induction.

For (i) write  $p = g' + (\text{LT}(p) + r')$

For (ii) write  $p = (g + g') + r'$

To argue uniqueness, suppose

$$p = g_1 + r_1 = g_2 + r_2 \quad \text{for } r_1 \neq r_2$$

Rearrange to get

$$r_1 - r_2 = g_2 - g_1 \in I$$

which shows that leading term in  $r_1 - r_2$   
is reducible. Contradiction □

The REDUCTION OPERATOR  $R_I$  is the operator that when applied to  $p$  returns the sum of irreducible terms  $R_I(p) = r$  such that  $p - r \in I$

Can think of  $r$  as representative of equivalence class of polynomials, or as "remainder" when dividing  $p$  by  $I$ .

(A bit like  $17 \bmod 5 = 2$ )

For set of (multilinear) polynomials  $P$ , write

$$\langle P \rangle = \{ \underline{q_i \cdot p_i} \mid p_i \in P, q_i \text{ polynomial} \}$$

for ideal generated by  $P$

In multilinear setting (or with Boolean axioms) we have

$$P \models q \iff q \in \langle P \rangle \iff R_{\langle P \rangle}(q) = 0$$

We won't prove this, because we don't need it, but it might be helpful for intuition.

Direction  $\Leftarrow$  is clear

Direction  $\Rightarrow$  needs work and uses Boolean axioms

Let us conclude our algebra recap with two more helpful facts

FACT B

For any two polynomials  $p, p'$  and ideals  $I_1 \subseteq I_2$ , it holds that  $R_{I_2}(p \cdot R_{I_1}(p')) = R_{I_2}(pp')$ .

ALG IV

This is the analogue of saying

$$\underline{a \cdot (b \bmod 15) \bmod 5 = ab \bmod 5}$$

Proof Write

$$p' = q' + r' \quad (1)$$

for  $q' \in I_1$ ,  $r'$  sum of irreducibles over  $I_1$ ,

$$p R_{I_1}(p') = pr' = q + r \quad (2)$$

for  $q \in I_2$ ,  $r$  sum of irreducibles over  $I_2$

Then combining (1) and (2) we get

$$pp' = pq' + pr' = pq' + q + r$$

where  $pq' + q \in I_2$  and  $r$  irreducible over  $I_2$ . By uniqueness (Fact A), get

$$R_{I_2}(pp') = r = R_{I_2}(p \cdot R_{I_1}(p')) \quad \square$$

FACT C

If  $t$  irreducible mod  $I$  and

$g: \text{Vars}(t) \rightarrow \mathbb{F}$  is any partial assignment s.t.

$t|_g \neq 0$ , then  $t|_g$  is also irreducible mod  $I$ .

*Set of irreducible monomials is downward-closed under restrictions.*

Proof Let  $t = m_g \cdot t'$  where  $m_g$  product of variables in  $\text{dom}(g)$  and by assumption  $\alpha = m_g|_g \neq 0$

Then  $t|_g = \alpha t'$ . If  $\exists q \in I$  s.t.  $\text{LT}(q) = t|_g$ , then  $\alpha^{-1} \cdot m_g \cdot q \in I$  and  $\text{LT}(\alpha^{-1} m_g q) = \alpha^{-1} m_g \cdot t|_g = m_g \cdot t' = t$ , contradicting that  $t$  is irreducible □